



UNIVERSITI PUTRA MALAYSIA

**MOLECULAR MARKERS AND THEIR APPLICATIONS IN
THE CONSTRUCTION OF GENETIC LINKAGE MAPS AND
ANALYSIS OF MONOGENIC AND QUANTITATIVE TRAITS
IN OIL PALM**

RAJINDER SINGH

FS 2005 36

**MOLECULAR MARKERS AND THEIR APPLICATIONS IN
THE CONSTRUCTION OF GENETIC LINKAGE MAPS AND
ANALYSIS OF MONOGENIC AND QUANTITATIVE TRAITS
IN OIL PALM**

By

RAJINDER SINGH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia
in Fulfilment of the Requirement for the Degree of Doctor of Philosophy**

July 2005



DEDICATION

To my parents, wife and children for their patience and love

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Doctorate of Philosophy

**MOLECULAR MARKERS AND THEIR APPLICATIONS IN THE
CONSTRUCTION OF GENETIC LINKAGE MAPS AND ANALYSIS OF
MONOGENIC AND QUANTITATIVE TRAITS IN OIL PALM**

By

Rajinder Singh

July 2005

Chairman: Professor Dr. Tan Soon Guan, PhD

Faculty: Faculty of Science

Molecular breeding is well suited to a perennial crop, like oil palm, in which the economic products are not produced until several years after planting. The use of DNA markers for selection in such crops could greatly reduce the number of breeding cycles needed. As such, the primary aim of this project was to map the oil palm genome by using restriction fragment length polymorphism (RFLP), amplified fragment length polymorphism (AFLP) and microsatellite (SSR) markers, in order to provide markers for selection in the breeding programme. A progeny derived from the selfing of a *tenera guineensis* palm (palm T128), which was segregating for the economically important monogenic characters of shell thickness and fruit color, was used for linkage map construction. A total of 523 segregating markers (117 RFLP, 22 SSR and 384 AFLP) were mapped to 17 linkage groups. In the current linkage map for the T128 palm, we have successfully mapped both the shell (*Sh*) gene and the fruit color (*Vir*) gene. Two RFLP markers mapped close to the *Vir* gene at a distance of 3 and 4 centiMorgans (cM), respectively. These two RFLP markers

were tested by using seven other independent crosses segregating for this trait. The markers could predict the trait with about 95% certainty. The *Sh* gene was located about 8 cM from the nearest marker (EAGG/MCTT-250, an AFLP marker). The AFLP marker could distinguish the *pisifera* palms (absence of shell) from the *dura* (thick shell) and *tenera* palms (thin shell) in the mapping family evaluated. These markers are useful tools for application in a molecular breeding programme for oil palm. Analysis was also extended to include mapping of quantitative traits (QTLs) associated with oil quality. Oil quality is determined by the fatty acid composition (FAC) in oil palm. By using the genome wide threshold levels calculated independently for each of the traits, significant QTLs were identified for myristic acid (C14:0), palmitic acid (C16:0), palmitoleic acid (C16:1) and stearic acid (C18:0). In the attempt at mapping additional QTLs associated with oil quality, an interspecific cross between a Colombian *oleifera* (UP1026) and the T128 palm was also examined. A map consisting of 412 markers (302 AFLP, 83 RFLP and 27 SSR) in 18 linkage groups was generated at a LOD score of 5.0. At a genome wide significance threshold of $P < 0.01$ and $P < 0.05$, significant QTLs were detected for iodine value (IV), C14:0, C16:0, C16:1 and C18:0. There were common markers revealing significant linkages for the same trait in both the mapping families analyzed. Such tagging of markers to economically important traits will aid in expediting the production of elite planting materials with greater precision. To facilitate oil palm genome analysis leading to physical mapping and the identification of molecular markers associated with QTL and map based cloning, we attempted to develop the tools and techniques needed to construct a bacterial artificial chromosome (BAC) library for oil palm. A suitable method to purify and prepare the single copy vector (pBeloBAC11) for BAC transformation was

established. The proper partial digestion conditions of oil palm megabase DNA for BAC library construction were also determined. Several oil palm BAC clones were successfully identified. Hybridization of these BAC clones with oil palm DNA as probe confirmed the presence of oil palm DNA in those clones. In map based cloning efforts, cosmid libraries (with cloning capacity of up to 50 kb) are useful tools to complement BAC libraries. For this reason we constructed a cosmid library for oil palm. The library contains 65,000 clones with insert size ranging from 30 kb to 42 kb. Hybridization of randomly selected cosmid clones with oil palm DNA also confirmed the presence of oil palm DNA in these clones.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Falsafah Kedoktoran

**PENANDA MOLEKUL DAN APLIKASINYA DALAM PEMBINAAN
RANGKAIAN GENETIK SERTA ANALISIS MONOGENIK DAN CIRI
KUANTITATIF DALAM KELAPA SAWIT**

Oleh

Rajinder Singh

Julai 2005

Pengerusi: Profesor Dr. Tan Soon Guan, PhD

Fakulti: Fakulti Sains

Biakbaka molekul amat bersesuaian dengan tumbuhan saka seperti kelapa sawit di mana penghasilan produk ekonomi tidak akan terhasil sehingga beberapa tahun selepas penanaman. Penggunaan penanda DNA (asid deoksiribonukleik) untuk pemilihan dalam tanaman seperti ini, dapat mengurangkan bilangan kitaran biakbaka yang diperlukan. Oleh itu, tujuan utama projek ini adalah untuk memetakan genom kelapa sawit dengan menggunakan teknik-teknik seperti RFLP ("restriction fragment length polymorphism"), AFLP ("amplified fragment length polymorphism") dan penanda mikrosatelit (SSR) dalam usaha menyediakan penanda untuk pemilihan dalam program biakbaka. Kacukan sendiri satu pokok tenera *guineensis* (pokok T128) yang didapati memberikan perbezaan ciri-ciri monogenik yang amat penting daripada segi ekonomi seperti ketebalan tempurung dan warna buah, telah digunakan untuk pemetaan genetik. Sejumlah 523 petanda informatif (117 RFLP, 22 SSR dan 384 AFLP) berjaya dipetakan kepada 17 kumpulan rangkaian. Pada peta rangkaian T128 tersebut, gen-gen berkaitan ketebalan

tempurung (*Sh*) dan warna buah (*Vir*) telah berjaya dipetakan. Dua penanda RFLP dipetakan berdekatan dengan gen *Vir* pada jarak 3 dan 4 centiMorgans (cM) masing-masing. Kedua-dua penanda RFLP tersebut telah diuji dengan menggunakan tujuh kacukan lain yang juga memberikan perbezaan pada ciri tersebut. Penanda-penanda tersebut boleh meramalkan 95% kehadiran ciri tersebut. Manakala gen *Sh* didapati berada 8 cM dari penanda yang berhampiran (EAGG/MCTT-250, penanda AFLP). Penanda AFLP ini berkeupayaan membezakan pokok *pisifera* (ketidakhadiran tempurung) daripada pokok *dura* (tempurung tebal) dan pokok *tenera* (tempurung nipis) dalam famili pemetaan yang diuji. Penanda-penanda ini amat berguna dalam aplikasi program biakbaka molekul kelapa sawit. Analisis juga dilanjutkan untuk memetakan lokus ciri-ciri kuantitatif (QTLs) yang berkaitan dengan kualiti minyak. Kualiti minyak kelapa sawit ditentukan oleh komposisi asid lemak (FAC). Dengan menggunakan tahap signifikasi genome yang dikira secara berasingan untuk setiap ciri, QTLs bererti dikenalpasti untuk asid miristik (C14:0), asid palmitik (C16:0), asid palmitoleik (C16:1) dan asid stearik (C18:0). Dalam percubaan pemetaan QTLs tambahan berkaitan dengan kualiti minyak, satu kacukkan interspesifik di antara *oleifera* Colombia (UP1026) dan T128 juga telah dikaji. Satu peta yang mengandungi 412 penanda (302 AFLP, 83 RFLP dan 27 SSR) dalam 18 kumpulan rangkaian telah dijana pada skor LOD bersamaan 5.0. Pada ambang signifikan genome $P < 0.01$ dan $P < 0.05$, QTL yang signifikan telah dikesan untuk nilai iodin (IV), C14:0, C16:0, C16:1 dan C18:0. Terdapat penanda yang sama mendedahkan rangkaian signifikan untuk ciri yang sama dalam kedua-dua keluarga pemetaan yang dikaji. Penandaan penanda terhadap ciri kepentingan ekonomi dapat membantu mempercepatkan penghasilan bahan tanaman elit dengan lebih jitu. Untuk membantu analisis genom kelapa sawit ke arah pemetaan fizikal dan

pengenalpastian penanda molekul yang berkaitan dengan QTL dan pengklonan berasaskan peta, kami telah membangunkan peralatan dan teknik yang diperlukan untuk membina perpustakaan kromosom bakteria buatan (BAC) kelapa sawit. Satu kaedah yang sesuai untuk menuliskan dan menyediakan vektor salinan tunggal (pBeloBAC11) untuk transformasi BAC telah diasaskan. Suatu keadaan pencernaan separa teratur DNA megabes kelapa sawit untuk pembinaan perpustakaan BAC telah ditentukan. Beberapa klon BAC kelapa sawit berjaya dikenalpasti. Penghibridan klon-klon BAC ini dengan DNA kelapa sawit sebagai prob mengesahkan kehadiran DNA kelapa sawit di dalam klon-klon tersebut. Dalam usaha pengklonan berasaskan peta, perpustakaan-perpustakaan kosmid (dengan kapasiti pengklonan sehingga 50 kilobes) merupakan asas yang berguna kepada komplementari perpustakaan BAC. Untuk tujuan ini, kami telah membina satu perpustakaan kosmid untuk kelapa sawit. Perpustakaan ini mengandungi 65, 000 klon dengan saiz selitan di antara 30 hingga 42 kb. Penghibridan beberapa klon kosmid terpilih dengan DNA kelapa sawit juga mengesahkan kehadiran DNA kelapa sawit di dalam klon-klon ini.

ACKNOWLEDGEMENTS

I would like to thank Prof. Dr. Tan Soon Guan, chairman of the supervisory committee for his patience, advice and guidance during this project. I would also like to extend my appreciation to Assoc. Prof. Dr. Jothi Malar Panandam, member of the supervisory committee, for her suggestions and valuable comments that helped towards the completion of this study. I would especially like to acknowledge Dr. Cheah Suan Choo, a colleague at MPOB and another member of the supervisory committee, without whom the work described in this study, would never have been initiated. Dr. Cheah's continuing faith in the utility of markers in oil palm breeding kept the work going and most importantly maintained funding for this work.

I would also like to extend my thanks to oil palm breeders especially, Dr. N. Rajanaidu (MPOB) and Mr. Mukesh Sharma (United Plantations), whose wholehearted support and cooperation allowed the work to be completed successfully. I would also like to thank the management at United Plantations for allowing their breeding material to be utilized in this study.

My heartfelt thanks and appreciation goes to my colleagues at MPOB, especially Ms. Rahimah Abd Rahman and En. Mohd Nor Ali for their sacrifices, patience and help throughout the study. My special thanks also goes to my laboratory colleagues Ms. Jayanthi Nagappan, Ms. Grace Ting Ngoot Chin, Mr. Chua Bok Hui, Mr. Leslie Low Eng Ti and Ms. Leslie Ooi Cheng Li for their support and encouragement throughout this study.

I certify that an Examination Committee met on 6th July 2005 to conduct the final examination of Rajinder Singh on his Doctor of Philosophy thesis entitled "Molecular Markers and their Applications in the Construction of Genetic Linkage Maps and Analysis of Monogenic and Quantitative Traits in Oil Palm" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Misri Kusnan, PhD

Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Siti Shapor Siraj, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Faridah Qamaruz Zaman, PhD

Lecturer
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Mikeal L. Roose, PhD

Professor
Department of Botany and Plant Science
University of California
United States of America
(External Examiner)



GULAM RUSUL RAHMAT ALI, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **22 AUG 2005**

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

Tan Soon Guan, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Jothi Malar Panandam, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

Cheah Suan Choo, PhD

Senior Research Fellow
Biology Division
Malaysian Palm Oil Board
(Member)



AINI IDERIS, PhD

Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 08 SEP 2005

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



RAJINDER SINGH

Date: 9/8/2005

TABLE OF CONTENTS

| | Page |
|---|--------|
| DEDICATION | 2 |
| ABSTRACT | 3 |
| ABSTRAK | 6 |
| ACKNOWLEDGEMENTS | 9 |
| APPROVAL | 10 |
| DECLARATION | 12 |
| LIST OF TABLES | 17 |
| LIST OF FIGURES | 20 |
| LIST OF ABBREVIATIONS | 25 |
| CHAPTER | |
| I. INTRODUCTION | 29 |
| II. LITERATURE REVIEW | |
| Oil Palm: Taxonomy, organ and distribution | 41 |
| The biology of oil palm | 42 |
| History of oil palm cultivation in Malaysia | 47 |
| Economic importance and future of oil palm | 53 |
| Products from oil palm | 57 |
| Biotechnology opportunities for oil palm | 59 |
| Molecular breeding in oil palm | 62 |
| DNA markers | 64 |
| i) Restriction Fragment Length Polymorphism (RFLP) markers | 66 |
| ii) Random Amplified Polymorphic DNA (RAPD) markers | 67 |
| iii) Amplified Fragment Length Polymorphism (AFLP) markers | 69 |
| iv) Microsatellite markers (Simple Sequence Repeats, SSR) | 72 |
| v) New marker systems | 76 |
| Application of DNA markers in genetic mapping | 77 |
| Using genetic mapping to study and improve agricultural productivity – molecular breeding | 85 |
| Success in marker assisted selection (MAS) or molecular breeding | 90 |
| Opportunities for DNA markers, genetic mapping and molecular breeding in oil palm | 93 |
| Genes influencing simply inherited traits | 95 |
| Genes influencing quantitative traits | 97 |
| Comparative mapping | 100 |
| Towards positional cloning (map based cloning) | 102 |

III. CONSTRUCTION OF A GENETIC LINKAGE MAP USING RFLP AND AFLP MARKERS AND MAPPING OF MONOGENIC AND QUANTITATIVE TRAITS IN OIL PALM

| | |
|---|-----|
| Introduction | 106 |
| Methodology | 114 |
| Plant materials | 114 |
| DNA extraction | 115 |
| Restriction enzyme (RE) digestion of DNA | 118 |
| Electrophoresis of genomic DNA for Southern analysis | 119 |
| RFLP probes | 120 |
| Labeling of selected DNA probes | 122 |
| Southern hybridization | 123 |
| AFLP analysis | 125 |
| i. <i>EcoRI/MseI</i> Enzyme Pair | 125 |
| ii. <i>TaqI/HindIII</i> | 127 |
| Digestion with <i>TaqI</i> and <i>HindIII</i> | 128 |
| Ligation to <i>TaqI</i> and <i>HindIII</i> Adapters | 128 |
| Pre-selective and selective amplification | 130 |
| Microsatellites | 132 |
| A. Isolation of Microsatellites in Oil Palm | 132 |
| B. Labeling of Microsatellite Primers and Analysis on Acrylamide Gel | 135 |
| C. Application of Published Oil Palm Microsatellite Sequences | 136 |
| Data analysis | 137 |
| Map construction | 138 |
| Evaluation of monogenic traits | 140 |
| Evaluation of quantitative traits | 141 |
| Results | 143 |
| DNA extraction | 143 |
| RFLP analysis | 144 |
| i. Selection of Informative Clones | 144 |
| ii) Inheritance Analysis of RFLP Loci in the Mapping Family. | 147 |
| AFLP analysis | 151 |
| Microsatellite (SSR) markers | 155 |
| i. Amplification Using Degenerate Primers | 155 |
| ii. Screening of Published Microsatellite Loci | 163 |
| Construction of genetic linkage map | 165 |
| Linkage of markers to shell gene and fruit color trait | 171 |
| Linkages of markers to quantitative traits | 180 |
| QTL analysis | 184 |
| Segregation of markers associated with QTLs | 188 |

| | |
|--|-----|
| Discussion | 191 |
| Choice of mapping families | 191 |
| DNA extraction | 192 |
| Marker analysis | 193 |
| Genetic map construction | 201 |
| Linkage to fruit color | 206 |
| Linkages to the shell gene locus | 207 |
| Linkage to quantitative traits (QTLs) | 209 |
| VI | |
| CONSTRUCTION OF GENETIC LINKAGE MAP AND ANALYSIS OF QUANTITATIVE TRAITS IN AN INTERSPECIFIC CROSS OF OIL PALM | |
| Introduction | 215 |
| Methodology | 220 |
| Plant materials | 220 |
| Preparation of genomic DNA | 220 |
| Amplified fragment length polymorphism (AFLP) procedure | 221 |
| Restriction fragment length polymorphism (RFLP) analysis | 222 |
| i. RFLP Probes | 222 |
| ii. Southern Blotting and Hybridization | 223 |
| Microsatellites | 224 |
| A. Isolation of Microsatellites in Oil Palm | 224 |
| B. Application of Published Oil Palm Microsatellite Sequences | 224 |
| Data analysis | 225 |
| Map construction | 226 |
| Quantitative data analysis | 227 |
| Results and Discussion | 229 |
| Mapping family | 229 |
| Screening a small sample size by using AFLP analysis | 230 |
| RFLP analysis | 234 |
| Microsatellite (SSR) markers | 238 |
| i) Screening of the Developed SSR Markers | 238 |
| ii) Screening of Published SSR Markers | 240 |
| Segregation analyses of AFLP, RFLP and SSR markers in the mapping family | 242 |
| i) AFLP Screening | 242 |
| ii) RFLP Screening | 250 |
| iii) SSR Markers | 251 |
| iv) Linkage analysis | 254 |
| Quantitative traits | 262 |
| QTL analysis | 269 |
| Segregation of markers associated with QTLs | 277 |
| Validation of marker-QTL linkage | 279 |

| | | |
|------------|--|-----|
| V | ISOLATION OF MEGABASE-SIZE DNA FROM OIL PALM AND ITS APPLICATION FOR CONSTRUCTION OF LARGE INSERT LIBRARIES | |
| | Introduction | 283 |
| | Methodology | 286 |
| I | BAC Library Construction | 286 |
| | Isolation of HMW DNA from Oil Palm Nuclei | 286 |
| | Restriction Digestion of HMW DNA | 287 |
| | Preparation of <i>Hind</i> III Partially Digested Oil Palm HMW DNA | 288 |
| | BAC Vector Preparation | 289 |
| | Ligation of DNA Fragments into the BAC Vector | 290 |
| | Transformation of Recombinant DNA Molecules (BAC) into <i>E.coli</i> by Electroporation | 291 |
| | Individual BAC Analysis | 291 |
| II. | COSMID Library Construction | 292 |
| | Preparation of Genomic DNA | 292 |
| | Ligation and Packaging of DNA | 293 |
| | Individual Cosmid Clone Analysis | 294 |
| | Results and Discussion | 295 |
| | Isolation of HMW DNA | 295 |
| | Partial digestion of HMW DNA | 299 |
| | Preparation of the pBeloBAC11 vector | 301 |
| | Construction and characterization of oil palm BAC clones | 303 |
| | Construction of cosmid library | 308 |
| | Preparation of genomic DNA for cosmid library construction | 311 |
| | Conclusion | 316 |
| VI | GENERAL DISCUSSION | 319 |
| VII | CONCLUSION | 328 |
| | REFERENCES | 337 |
| | APPENDICES | 364 |
| | BIODATA OF AUTHOR | 423 |

List of Tables

| Tables | Page |
|--|------|
| 2.1 Comparison of <i>dura</i> , <i>tenera</i> and <i>pisifera</i> fruit forms | 49 |
| 2.2 World production of 17 oils and fats (2000-2003) ('000 Tonnes) | 54 |
| 2.3 Comparative yields for vegetative oils (1995-1996) | 55 |
| 2.4 Some naturally occurring fatty acids in plants | 58 |
| 2.5 Fatty acid composition (%) of palm oil and palm kernel oil | 59 |
| 2.6 Application of DNA markers | 64 |
| 2.7 Comparison of marker systems | 65 |
| 2.8 Informativeness of some common mapping families | 83 |
| 3.1 Below illustrates the 2 different types of segregation patterns observed in the mapping family from RFLP and AFLP markers | 138 |
| 3.2 Source of cDNA probes screened for RFLP | 145 |
| 3.3 Number and proportion of probes revealing RFLPs in the mapping family | 147 |
| 3.4 Summary of RFLP, AFLP and SSR analyses of the 136 progeny of the mapping family | 150 |
| 3.5 The total number of bands and segregating markers scored using 69 AFLP primer pairs in the 136 progeny of the mapping family | 153 |
| 3.6 No. of recombinant colonies produced using two different primers PCT1 and PCT4 | 157 |
| 3.7 Microsatellites isolated with the 5' anchor PCR method. N is the number of clones sequenced for each anchor primer | 158 |
| 3.8 Microsatellite primer sets, primer sequences, and optimum annealing temperatures | 160 |
| 3.9 Published oil palm SSR primer set used to analyze the mapping family | 164 |
| 3.10 Analysis of palms for fruit type and color | 172 |
| 3.11 Correlation of <i>virescens</i> and <i>nigrescens</i> palms with expected banding profile for probes MET16 and KT3 | 177 |



| | | |
|------|--|-----|
| 3.12 | Linkage of RFLP markers, MET16 and KT3 to the fruit color locus in seven different families. | 180 |
| 3.13 | Means of the male parent (Palm T128) and F ₁ progeny obtained for carotene content, IV and fatty acid composition. | 181 |
| 3.14 | Normality assessment carried out using the Shapiro-Wilk test | 183 |
| 3.15 | Correlation between fatty acids (n = 97) in the F ₁ progeny. | 184 |
| 3.16 | QTLs for fatty acid composition found to be significant at the empirical genome wide mapping threshold. | 186 |
| 3.17 | Correlation between segregation of markers linked to QTLs and specific traits. | 190 |
| 4.1 | Fatty acid composition of oil from various sources of <i>E. oleifera</i> germplasm. | 216 |
| 4.2 | Segregation of AFLP amplification products in the progeny of the UP1026 (<i>E. oleifera</i>) x T128 (<i>E. guineensis</i>) interspecific cross. | 226 |
| 4.3 | Segregation of AFLP markers in the pseudo-testcross revealed by a subset of 26 primer pairs used in analysis of the <i>oleifera</i> x <i>guineensis</i> progeny. | 232 |
| 4.4 | Polymorphisms detected by cDNA probes in the interspecific hybrid mapping family. | 235 |
| 4.5 | Segregation of RFLP probes in the parental gametes. | 237 |
| 4.6A | Microsatellite primer sets, primer sequences, and optimum annealing temperatures. | 238 |
| 4.6B | Published microsatellite primer sets and primer sequences tested on the two mapping family. | 241 |
| 4.7 | Segregation of AFLP markers in the <i>oleifera</i> x <i>guineensis</i> progeny. | 246 |
| 4.8 | Summary of RFLP, SSR and AFLP analysis of the interspecific hybrid mapping family. | 249 |
| 4.9 | Means of the male parent (T128), female parent (UP1026) and F ₁ progeny obtained for carotene content, IV and fatty acid composition. | 264 |
| 4.10 | Assessment of Normality for the various traits using the Shapiro-Wilk test | 268 |

| | | |
|------|---|-----|
| 4.11 | Correlation between fatty acids (n = 81) in F ₁ progeny. | 269 |
| 4.12 | QTLs for IV and fatty acid composition found to be significant at the empirical genome wide mapping threshold. | 270 |
| 4.13 | QTLs for IV and fatty acid composition found to be significant at the empirical chromosome wide mapping threshold. | 276 |
| 4.14 | Correlation between segregation of markers linked to QTLs and specific traits. | 279 |
| 4.15 | Comparison of common markers influencing QTLs detected in both mapping families. | 281 |
| 5.1 | The three methods tested for pBeloBAC11 purification. The quality of the vector was assessed by examining an aliquot on a 1% TAE gel. | 302 |
| 5.2 | Analysis of transformation efficiency of BAC clones. | 305 |

List of Figures

| Figure | Page |
|---|------|
| 2.1 The oil palm fruit forms. | 45 |
| 2.2 Crossing strategies for developing RILs (recombinant inbred lines) and BCs (backcrosses). | 82 |
| 3.1 Ethidium bromide stained agarose gels showing typical profile obtained during trial digestion of seven independent DNA samples. Lane 1 is undigested DNA Lane 2 and 3 represent DNA digested with <i>EcoRI</i> and <i>HaeIII</i> , respectively. | 144 |
| 3.2 Relationship between the number of restriction enzymes used and the percentage of probes revealing RFLP. | 146 |
| 3.3 Segregation profile of probe SFB138 and SFB62 on a small sample of the progeny of the mapping family. The probe SFB 138 (as in A above) shows 1:2:1 segregation profile while probe SFB62 (as in B above) shows 3:1 segregation profile. "P" is the parental palm, T128. Arrows (\hookrightarrow) indicate the segregating bands. | 148 |
| 3.4 Hybridization of probe SFB 47 on a small sample of the progeny of the mapping family. Bands not present in the parental palm are indicated by an arrow (\blackrightarrow). The parental palm lane is indicated with the alphabet "P". Palms having such additional bands were excluded from the analysis of the mapping family. | 149 |
| 3.5 Duplicated loci observed with probe MT40 (A) and SFB56 (B). "P" is the parental palm. | 151 |
| 3.6 Segregation of AFLP loci in the progeny of a selfed tenera (T128) for the primer combinations, E-ACT/M-CAG (A) and T-AGG/H-CAT (B). "T128" refers to the parental palm. Black arrows indicate segregating bands. | 155 |
| 3.7 Amplification of oil palm genomic DNA, Nigerian <i>guineensis</i> (palm, T128) and Colombian <i>oleifera</i> (UP1026), using the anchor primers, PCT4 and PCT1. | 156 |
| 3.8 <i>EcoRI</i> digestion of clones randomly selected from the microsatellite enriched library. Only clones with insert sizes above 100bp were sequenced. M is 100 bp ladder. | 157 |
| 3.9 Determination of annealing temperature of the primer pairs designed flanking the microsatellite motifs. The primer pairs were used to amplify oil palm genomic DNA (parental palm, T128) at the appropriate annealing temperature (ranging from 46-56 °C) and subsequently analyzed on a 1% agarose gel. M is 100 bp ladder. | 159 |



| | | |
|--------|---|-----|
| 3.10 | Autoradiogram obtained with four different SSR primers. Parental palm, T128, is indicated with alphabet 'P'. Panel A (P1O14a) and Panel B (P4T8) are part of the SSR primers developed during this study. Panel C (CIR8) and Panel D (CNH0887) were part of the published SSR primer sets used in this study. Panel B also shows the duplicated loci observed. | 162 |
| 3.11 | Combined AFLP, RFLP and SSR map of T128 selfed. | 167 |
| 3.12 | Comparison of the closest marker identified to the shell gene (EGC/MCTT-250) with the phenotype scoring in the mapping population. | 173 |
| 3.13A | Segregation of the marker MET16 compared to that of the fruit color gene in the mapping population. N is <i>nigrescens</i> and V is <i>virescens</i> fruits. Symbol '*' indicates fruit color not matching banding profile. | 175 |
| 3.13B: | Segregation of the marker KT3 compared to that of the fruit color gene in the mapping population. N is <i>nigrescens</i> and V is <i>virescens</i> fruits. Symbol '*' indicates fruit color not matching banding profile. | 176 |
| 3.14A | Screening of the marker MET16 linked to the fruit color locus, on seven independent populations. V and N refer to <i>virescens</i> and <i>nigrescens</i> fruits, respectively. Arrows indicate segregation of the probe in these populations distinguishing the <i>virescens</i> and <i>nigrescens</i> fruits. The symbol '*' denotes palms showing a mismatch with the expected profile due to recombination between the marker and the fruit color loci. | 178 |
| 3.14B | Autoradiographs generated by screening the marker KT3 (also linked to the fruit color locus) on the same seven independent populations. V and N refer to <i>virescens</i> and <i>nigrescens</i> fruits, respectively. Arrows indicate segregation of the probe in these populations distinguishing the <i>virescens</i> and <i>nigrescens</i> fruits. The symbol '*' denotes palms showing a mismatch with the expected profile due to recombination between the marker and the fruit color loci. | 179 |
| 3.15 | Frequency distribution for carotene content, IV and fatty acid Components (C14:0, C16:0, C16:1, C18:0, C18:1, C18:2 and C18:3) in the F ₁ progeny. | 183 |
| 3.16 | QTL graphs for fatty acid composition (C14:0, C16:0, C16:1 and C18:0). | 186 |
| 4.1 | Segregation of AFLP loci in the progeny of the interspecific cross for the primer combinations EACT/MCAT (A) and TACG/HCTA (B). (←) indicates segregating bands. | 233 |

| | | |
|-----|---|-----|
| 4.2 | Autoradiograms showing the segregating loci revealed by RFLP markers KT35 (A), KT19 (B) and FDA39 (C). UP1026 and T128 are the female parent, Colombian <i>oleifera</i> and male parent, Nigerian <i>guineensis</i> , respectively. (←) indicates segregating bands. | 236 |
| 4.3 | Contaminating palm (marked X) detected in interspecific cross with primer pair E-ACC/M-CAC. P1 and P2 refer to parents UP1026 and T128, respectively. | 244 |
| 4.4 | Microsatellite profiles in a small subset of the interspecific cross (UP1026xT128). The profiles shown were obtained using primer P4T8 (A), P1T6b (B), CIR377(C) and CNH0887(D). | 253 |
| 4.5 | Combined AFLP, RFLP and SSR map of interspecific hybrid (palm T128) | 257 |
| 4.6 | Frequency distribution for carotene content, IV and fatty acid components (C14:0, C16:0, C16:1, C18:0, C18:1, C18:2 and C18:3) in the interspecific hybrid F ₁ progeny. | 267 |
| 4.7 | QTL graphs for IV and fatty acid composition (C14:0, C16:0, C16:1, C18:0, C18:1, C18:2). | 272 |
| 4.8 | Comparison of Group 10 (from the interspecific cross) and Group 12 (from the T128 selfed cross, from Chapter three), showing common RFLP (showed by line) and AFLP (dotted line) markers mapped. The common loci influencing the QTLs for C14:0 and C16:1 are also indicated. The bar shows the range for QTL obtained from Interval Mapping while ovoid shapes indicate the markers identified through the Kruskal-Wallis test to be influencing the QTLs concerned. | 282 |
| 5.1 | PFG Electrophoresis of megabase DNA prepared from oil palm. The gel (1% agarose, 0.5 x TBE) was run with a CHEF-DR111 apparatus (Biorad) at 6 V/cm, 14 °C and ramping times of 80 s and 100 s for 20 hours. Lanes 1 to 6 indicate megabase DNA prepared in agarose plugs. M is yeast chromosome size standards. | 296 |
| 5.2 | Gel analysis of purified megabase DNA from oil palm. Lanes 1-2 show unpurified DNA, Lanes 3-5 show purified DNA. Lane M is yeast chromosome used as a size marker and M1 is Lambda ladder PFG marker. | 297 |
| 5.3 | (A) Digestion of megabase DNA with restriction enzymes as indicated. N: <i>NotI</i> ; E: <i>EcoRI</i> , B: <i>BamHI</i> and U: Undigested. Lane M is lambda ladder PFG marker. (B): Southern hybridization of Gel in A hybridized with probe G-142 (single copy probe from oil palm). | 299 |

- 5.4 Partial digestion of oil palm DNA. The megabase-size DNA in agarose plugs was partially digested with various concentrations of *HindIII* as indicated at the top of each lane. Lane M is PFG lambda ladder size marker. 300
- 5.5 DNA fragments purified from low melt agarose gel and examined on a Pulse Field Gel. Lane 1: 50-100kb, Lane 2: 100-200 kb, Lane 3: 200-250 kb and Lane M: PFG lambda ladder. 301
- 5.6 Preparation of pBeloBAC11 plasmid for ligation with genomic DNA. Lane 1 indicates plasmid digested by the cloning enzyme *HindIII*. Lane 2: Plasmid digested (*HindIII*), dephosphorylated and treated with T4 DNA ligase. Lane 3: Plasmid digested (*HindIII*), dephosphorylated, treated with kinase (to restore 5' phosphate group) and subsequently with T4 DNA ligase. Lane M is 1kb ladder (Gibco BRL, USA). 303
- 5.7 Analysis of oil palm BAC clones by PFG electrophoresis. (A) Ethidium bromide stained CHEF gel. Lanes M1 and M are Mid-Range PFG marker and PFG lambda ladder, respectively (NEB, USA). Lanes 3-13, alkaline lysis mini preparations of randomly picked recombinant BAC clones derived from size-selected DNA and digested with *NotI*. (B) Southern hybridization of gel in A hybridized with radioactively labeled total oil palm DNA. 306
- 5.8 Preparation of *SuperCos1* cosmid vector. Lane 1 is undigested plasmid. Lane 2 is plasmid digested with *XbaI* producing as expected a single band at 7.3 kb. Lane 2 is the plasmid further digested with *BamHI*, to produce the 2 vector arms at 1.1kb and 6.3 kb. 309
- 5.9 Analysis of the dephosphorylation of the *SuperCos1* cosmid vector after *XbaI* digestion and prior to *BamHI* digestion. Lane 1 is undigested plasmid. Lane 2 is plasmid digested with *XbaI*, dephosphorylated and treated with T4 DNA Ligase. Lane 3 is plasmid digested with *XbaI*, dephosphorylated, treated with T4 polynucleotide kinase (to restore 5' phosphate group) and subsequently with T4 DNA ligase. 310
- 5.10 PFG electrophoretic analysis of genomic DNA to be used for cosmid library construction. The DNA was analysed with a CHEFDR11 system on a 1% agarose gel in 0.5 x TBE at 14 °C at 6.0 V/cm for 16 hours with pulse time of 20 s. Lane 1 and 2 is D x P oil palm DNA, isolated using the traditional CTAB method. Lane 3 is PFG lambda ladder (NEB, USA). 311

- 5.11 Analysis of genomic DNA used in cosmid library construction. Panel A shows analysis of digestion on a 0.3% (TAE) agarose gel. Lane 1: Lambda-DNA *Hind*III digest (NEB, USA), Lane 2: undigested DNA, Lane 3-5: DNA digested with *Sau*3A1 at 0.04 U, 0.06 U and 0.08 U/ μ g DNA, respectively. Panel B is analysis of digestion on a PFG electrophoresis. Lane 1: undigested DNA, Lane 2-5 is DNA digested with 0.04 U/ μ g DNA. Lane M is Lambda Ladder PFG marker (NEB, USA). PFG was run on a 1% agarose gel (0.5 x TBE) at 4.5 V/cm for 16 hours at 10 s pulse time. 312
- 5.12 Analysis of oil palm cosmid clones on a 0.8% TAE agarose gel. Panel A (lane 2-9) shows alkaline lysis preparation of randomly picked cosmid clones digested with *Eco*R1. Panel B is Southern hybridization of gel in A hybridized with radioactively labeled total oil palm DNA. Lane 1 is Lambda-DNA *Hind*III digested (NEB, USA). 314